

WiQo BALANCING CREAM

HOME CARE

NEW



SEBUM-REGULATING

OILY SKIN

ACNE-PRONE SKIN

WiQo Balancing Cream

The **new WiQo sebum-regulating and balancing cream** has a dry texture that is perfect for impure skins.

Thanks to the **synergistic formulation** of its active principles, it is indicated for skins with excess sebum, blackheads, dilated pores and acne tendency.



CLINICALLY PROVEN RESULTS

PORE VOLUME
-38.3%

Improved subjects
100%

NUMBER OF PORES
-32.8%

Improved subjects
95%

SKIN SEBUM
-40.7%

Improved subjects
95%

INSTRUCTIONS FOR USE

Cleanse the skin accurately.

Apply **WiQo Balancing Cream** on the entire face once or twice a day. The cream should be used for at least 2 months.

ACTIVE INGREDIENTS

COMEDOLYTIC

- SALICYLIC ACID
- AZELAIC ACID

SEBUM SECRETION REGULATION

- AZELAIC ACID (20%)
- BIFIDA FERMENT LYSATE
- FERMENTED FOLIC ACID

SOOTHING

- ADA
- FERMENTED CAMELLIA OIL
- PHYTOSPHINGOSINE

VITAMIN E

FERMENTED EXTRACTS

NIACINAMIDE

- BIFIDA FERMENT LYSATE
- FERMENTED FOLIC ACID
- FERMENTED CAMELLIA OIL

INGREDIENTS: AQUA (Water), AZELAIC ACID, BIFIDA FERMENT LYSATE, ETHOXYDIGLYCOL, AZELAMIDOPROPYL DIMETHYL AMINE, SALICYLIC ACID, POLYGLYCERYL-2 STEARATE, GLYCERIN, PROPANEDIOL, GLYCERYL STEARATES, STEARYL ALCOHOL, STEARYL LINOLEATE, BUTYLENE GLYCOL, CAPRYLIC/CAPRIC TRIGLYCERIDE, PSEUDOSZYMA EPICOLA/CAMELIA JAPAONICA SEED OIL FERMENT EXTRACT FILTRATE, SODIUM ACRYLATE/SODIUM ACRYLOYLDIMETHYL TAURATE COPOLYMER, DIMETHICONE, PHENOXYETHANOL, DEXTRIN, OCTYLDODECANOL, STEARYL STEARATE, BACILLUS/FOLIC ACID FERMENT FILTRATE EXTRACT, SODIUM GLUCURONATE, POLYDEXTROSE, AMYLOPECTIN, NIACINAMIDE, OCTYLDODECYL OLEATE, PHYTOSPHINGOSINE, TRIDECETH-6, PEG/PPG-18/18 DIMETHICONE, POLYHYDROXYSTEARIC ACID, ETHYLHEXYLGLYCERIN, TOCOPHEROL, OCTYLDODECYL STEAROYL STEARATE, SODIUM HYALURONATE, 1,2-HEXANEDIOL, HELIANTHUS ANNUUS SEED OIL (Helianthus Annuus (Sunflower) Seed Oil), XANTHAN GUM.

BEFORE AND AFTER

Clinical tests* have demonstrated the sebum-regulating and pore-reducing efficacy of the **WiQo sebum-regulating and balancing cream**. The texture of the epidermis is visibly improved after only a short time of use.

* Evaluation of sebum-regulating and pore reduction action. *In vivo* test on 20 volunteers with mild to moderate acne (I-III degree). Product applied twice a day, in the morning and in the evening, for 4 weeks.




Before



After 4 weeks °



Before



After 4 weeks °

° **WiQo Balancing Cream** applied twice a day, in the morning and in the evening, for 4 weeks

BIBLIOGRAPHY

- Akarsu et al. Journal of Dermatology 2012; 39 433-8
Efficacy of the addition of salicylic acid to clindamycin and benzoyl peroxide combination for acne vulgaris
- Borelli et al Acta Derm Venereol 2006; 86 316-319
In vivo Porphyrin Production by P. acnes in Untreated Acne Patients and its Modulation by Acne Treatment
- J Biol Chem. 2014 Oct 17; 289(42) 28846–28855
RecO Protein Initiates DNA Recombination and Strand Annealing through Two Alternative DNA Binding Mechanisms
- Lee et al. Dermatologic Surgery 2003; 29(12) 1196-9
Salicylic Acid Peels for the Treatment of Acne Vulgaris in Asian Patients
- Liu H et al. Cochrane Database of Systematic Reviews 2020, Issue 5
Topical azelaic acid, salicylic acid, nicotinamide, sulphur, zinc and fruit acid (alpha-hydroxy acid) for acne (Review)
- Liu et al. Cochrane Database of Systematic Reviews 2014
Topical azelaic acid, salicylic acid, nicotinamide, and sulphur for acne (Protocol)
- M.Shu et al Curr Med Chem 2013 20 (4) 562-568
Porphyrin Metabolisms in Human Skin Commensal Propionibacterium acnes Bacteria: Potential Application to Monitor Human Radiation Risk
- Kim.S Et Al, Mol. Med 12 (1 - 3) 17 - 24 , January, 2006
Phytosphingosine Stimulates the Differentiation of Human Keratinocytes and Inhibits TPA-Induced Inflammatory Epidermal Hyperplasia in Hairless Mouse Skin
- S. Puebla-Barragan,G. Reid Molecules-26-01249
Probiotics in Cosmetic and Personal Care Products: Trends and Challenges
- JQ. Del Rosso et. Al. Cutis. 2011;8867-72
Azelaic Acid Gel 15% in the Management of Papulopustular Rosacea: A Status Report on Available Efficacy Data and Clinical Application
- Tamara Searle et al., Journal of Dermatological Treatment. 2020 1-31
The versatility of azelaic acid in dermatology



For information: scientificdivision@wiqo.eu

GPQ srl

Operational headquarters: Via Martinelli, 16 - 34015 Muggia (Trieste) - Italy
Tel.: +39 040 360408 - Fax: +39 040 761393 - gpq@wiqo.eu
www.wiqo.com